



SPECIFICATIONS



AgriBusiness[®]
ANIMAL NUTRITION

© Perdue AgriBusiness, LLC.
All rights reserved.
Perdue[®] is a registered
trademark of Perdue, Inc.
6906 Zion Church Rd.
Salisbury, MD 21804

ProvAAI² and Metaatein[®]
are registered trademarks
of Perdue, Inc.

Standard Analysis	% DM
Moisture	7.8
Dry Matter	92.2
Nitrogen	12.9
Crude Protein	80.5
Soluble Protein	25.2
Ammonia	—
NDIP	13.6
ADIP	13.6
Crude Fiber	1.6
ADF	2.4
NDF	5.9
Lignin	0.4
Sugar	0.9
Starch	0.4
Soluble Fiber	1.3
NFC	2.5

Protein Bio-Availability	% DM
RUP % CP	90.6
RUPd % RUP	81.4
dRUP % DM	59.4
dRUP-Lys % DM	4.5
dRUP-Meth % DM	0.6
dRUP-EAA % DM	31.1

Macro Mineral	% DM
Ash	7.0
Calcium	1.5
Phosphorus	1.0
Magnesium	0.1
Potassium	0.5
Sulfur	0.8
Sodium	0.5
Chloride	0.3

Trace Mineral	ppm
Iron	1468.2
Manganese	13.0
Zinc	58.7
Copper	10.8

Fatty Acids	% DM
Ether Extract	4.1
Total Fatty Acids	3.8
Glycerol	0.2
Pigment	0.1
C12:0	—
C14:0	—
C16:0	0.9
C16:1	0.1
C18:0	0.4
C18:1T	—
C18:1C	1.2
C18:2	1.0
C18:3	—
Other	0.2
Fat Type	2.0
Lipolysis Rate	137.0

Pool	% DM	Rate	Int dig
CHO A1	—	—	—
CHO A4	0.92	40.00	100.0
CHO B1	0.36	17.00	75.0
CHO B2	1.26	30.00	75.0
CHO B3	4.81	7.00	20.0
CHO C	1.06	—	—
PRO A1	—	—	—
PRO A2	25.21	1.51	100.0
PRO B1	41.74	0.91	100.0
PRO B2	—	—	—
PRO C	13.55	—	—

Amino Acids	Product		Model Specifications ¹	
	% DM	% of CP	% DM	% of CP
Arginine	3.8	4.7	3.6	4.4
Histidine	4.0	5.0	4.6	5.7
Isoleucine	1.3	1.7	1.1	1.4
Leucine	9.5	11.8	9.9	12.3
Lysine	5.6	7.0	6.1	7.6
Methionine	0.9	1.1	0.8	1.0
Phenylalanine	5.0	6.2	5.2	6.5
Threonine	3.3	4.1	3.3	4.1
Tryptophan	0.9	1.1	0.9	1.1
Valine	6.4	8.0	6.6	8.2
EAA	40.7	50.6	42.1	52.3

¹When a feed component of a dairy ration consists of a blend of feed ingredients differing in rumen undegradable protein and/or post-ruminal protein digestibility, the amino acids of the resulting mixture differ in their ruminal degradability and post ruminal digestibility. Most ration formulation software allow only a single, common set of parameters to characterize ruminal degradability and post ruminal digestibility of all essential amino acids; one cannot enter parameters specific to each amino acid. Therefore, entering the amino acid profile of the whole protein of a feed mixture results in erroneous values of its metabolizable amino acid content. To circumvent this problem, users should use the 'model specifications' amino acid values in software such as CNCPs/NRC so that the correct supply of each metabolizable amino acid is calculated correctly.

Science based. Research driven.®