

SPECTRUM



SPECIFICATIONS



AgriBusiness®
ANIMAL NUTRITION

© Perdue AgriBusiness, LLC.
All rights reserved.
Perdue® is a registered trademark of Perdue, Inc.
6906 Zion Church Rd.
Salisbury, MD 21804
SPECTRUM AgriBlue
is a registered trademark of Perdue, Inc.

Standard Analysis	% DM
Moisture	6.5
Dry Matter	93.5
Nitrogen	14.7
Crude Protein	92.1
Soluble Protein	41.3
Ammonia	—
NDIP	7.8
ADIP	7.8
Crude Fiber	—
ADF	—
NDF	—
Lignin	—
Sugar	—
Starch	—
Soluble Fiber	—
NFC	—

Fatty Acids	% DM
Ether Extract	2.6
Total Fatty Acids	2.4
Glycerol	0.1
Pigment	0.1
C12:0	—
C14:0	—
C16:0	0.4
C16:1	0.1
C18:0	0.7
C18:1T	—
C18:1C	0.6
C18:2	0.5
C18:3	—
Other	0.1
Fat Type	1.8
Lipolysis Rate	55.7

Protein Bio-Availability	% DM
RUP % CP	92.7
RUPd % RUP	90.9
dRUP % DM	77.6
dRUP-Lys % DM	6.7
dRUP-Meth % DM	4.3
dRUP-EAA % DM	45.1

Pool	% DM	Rate	Int dig
CHO A1	—	—	—
CHO A4	—	—	—
CHO B1	—	—	—
CHO B2	—	—	—
CHO B3	—	—	—
CHO C	—	—	—
PRO A1	—	—	—
PRO A2	41.25	1.08	100.0
PRO B1	43.06	0.60	100.0
PRO B2	—	—	—
PRO C	7.78	—	—

Macro Mineral	% DM
Ash	5.3
Calcium	0.7
Phosphorus	0.4
Magnesium	0.1
Potassium	0.3
Sulfur	0.7
Sodium	0.5
Chloride	0.5

Trace Mineral	ppm
Iron	2326.6
Manganese	7.3
Zinc	36.2
Copper	7.5

Amino Acids	Product		Model Specifications ¹	
	% DM	% of CP	% DM	% of CP
Arginine	3.7	4.1	3.6	3.9
Histidine	5.1	5.6	5.4	5.9
Isoleucine	0.9	1.0	0.8	0.9
Leucine	11.3	12.3	11.5	12.5
Lysine	7.7	8.4	7.9	8.6
Methionine	5.0	5.4	5.1	5.5
Phenylalanine	6.2	6.7	6.3	6.9
Threonine	4.0	4.3	3.9	4.3
Tryptophan	1.1	1.2	1.1	1.2
Valine	7.8	8.4	7.8	8.5
EAA	52.9	57.4	53.5	58.1

¹When a feed component of a dairy ration consists of a blend of feed ingredients differing in rumen undegradable protein and/or post-ruminal protein digestibility, the amino acids of the resulting mixture differ in their ruminal degradability and post-ruminal digestibility. Most ration formulation software allow only a single, common set of parameters to characterize ruminal degradability and post-ruminal digestibility of all essential amino acids; one cannot enter parameters specific to each amino acid. Therefore, entering the amino acid profile of the whole protein of a feed mixture results in erroneous values of its metabolizable amino acid content. To circumvent this problem, users should use the 'model specifications' amino acid values in software such as CNCPS/NRC so that the correct supply of each metabolizable amino acid is calculated correctly.

Science based. Research driven.®